



WORLD HEALTH ORGANIZATION
ORGANISATION MONDIALE DE LA SANTE

25649

DISTR.: LIMITED
DISTR.: LIMITEE
WHO/TB/88.154
ENGLISH ONLY

ANNUAL RISK OF TUBERCULOUS INFECTION

by

G.M. Cauthen¹, A. Pio² and H.G. ten Dam³



- 1
Division of Tuberculosis Control, U.S. Centers for Disease Control, Atlanta, Georgia, USA.
- 2
Formerly Chief, Tuberculosis and Respiratory Infections Unit, Division of Communicable Diseases, World Health Organization, Geneva, Switzerland.
- 3
Scientist, Tuberculosis Unit, Division of Communicable Diseases, World Health Organization, Geneva, Switzerland.

This document is not issued to the general public, and all rights are reserved by the World Health Organization (WHO). The document may not be reviewed, abstracted, quoted, reproduced or translated, in part or in whole, without the prior written permission of WHO. No part of this document may be stored in a retrieval system or transmitted in any form or by any means - electronic, mechanical or other without the prior written permission of WHO.

The views expressed in documents by named authors are solely the responsibility of those authors.

Ce document n'est pas destiné à être distribué au grand public et tous les droits y afférents sont réservés par l'Organisation mondiale de la Santé (OMS). Il ne peut être commenté, résumé, cité, reproduit ou traduit, partiellement ou en totalité, sans une autorisation préalable écrite de l'OMS. Aucune partie ne doit être chargée dans un système de recherche documentaire ou diffusée sous quelque forme ou par quelque moyen que ce soit - électronique, mécanique, ou autre - sans une autorisation préalable écrite de l'OMS.

Les opinions exprimées dans les documents par des auteurs cités nommément n'engagent que lesdits auteurs.

TABLE OF CONTENTS

	<u>Page</u>
I. INTRODUCTION	3
II. METHODS	3
III. RESULTS	5
IV. DISCUSSION	8
V. CONCLUSIONS	10
REFERENCES	11

FIGURES:

FIGURE 1: Recent estimates of annual risk of tuberculosis infection in developing countries	19
FIGURE 2: Annual risk of tuberculosis infection I. WHO African Region (except Algeria)	20
FIGURE 2: (continued) Annual risk of tuberculosis infection II. WHO Region of the Americas	21
FIGURE 2: (continued) Annual risk of tuberculosis infection III. WHO South-East Asia region	22
FIGURE 2: (continued) Annual risk of tuberculosis infection IV. WHO Eastern Mediterranean Region (and Algeria)	23
FIGURE 2: (continued) Annual risk of tuberculosis infection V. WHO Western Pacific Region	24

TABLES:

TABLE 1. ANNUAL RISK OF TUBERCULOSIS INFECTION IN DEVELOPING COUNTRIES BASED ON TUBERCULIN SURVEYS:	
I. WHO African Region (except Algeria)	26
II. WHO Region of the Americas	28
III. WHO South-East Asia Region	29
IV. WHO Eastern Mediterranean Region (and Algeria)	31
V. WHO Western Pacific Region	34

I. INTRODUCTION

Risk of infection in a population is in many respects the most informative index of the magnitude of the tuberculosis problem (Sutherland, 1976; Pio, 1984). The risk of infection at a particular time indicates the current magnitude of the incidence and prevalence of infectious cases (Styblo, 1985) and also indicates the magnitude of the tuberculosis problem years into the future. An observed decline in the risk of infection would be the earliest indicator of a decline in the epidemic cycle of tuberculosis, resulting from tuberculosis control activities or improvements in living standards. A rising risk of infection would be an early indicator of changes in the other direction, signalling the introduction of new risk factors, such as the spread of human immunodeficiency virus (HIV) infection.

It is known that risk of infection has been declining for many years in developed countries, but remains at high levels in many developing countries (Styblo, 1984). In developing countries in the last decade, a number of surveys of infection prevalence have been carried out in national populations as well as in smaller populations.

Therefore a project was carried out to assess the current level and trend in the risk of infection in developing countries by reviewing and assembling tuberculin skin test survey data available since 1975.

II. METHODS

Tuberculin skin test data collected since 1975 for populations in developing countries were assembled from reports to the World Health Organization and from the published literature. Data from surveys of childhood age groups that were judged of sufficient quality were selected in order to provide as valid and up to date an assessment as practical of the magnitude of the risk of infection. Comparable prior data for the same countries were also selected in order to judge whether the risk of infection is likely to have declined.

Prevalence of infection observed in childhood age groups was used to derive the average annual risk of infection that would have resulted in the observed cumulative prevalence rate. Choosing younger age groups allows the calculated average annual risk to be bracketed within a relatively narrow period of time between the average birth date of the group and the date of the survey (Styblo et al, 1969).

Each survey was judged on the basis of available documentation for ability to represent the population and to detect the proportion infected at a particular time. A sample survey was judged to have measured the proportion infected in the target population if the probability sampling design and the estimation method appeared correct in concept and conduct, and skin testing technique appeared adequate to measure the proportion infected in the sampling units.

Basis for selection

Surveys were selected which met most of the following criteria:

Specified the sampling design including the sampling frame, staging, stratification, sampling units, allocation, sampling weights, and estimation formulas and documentation to support that the design was followed with adequate coverage.

Provided reaction-size distributions for surveyed age groups and for bacteriologically confirmed cases from the same population in order to assess the definition of infection and in order to judge technique.

Documented type and strength of antigen used, technique of administration and reading, and quality control procedures.

Specified the methods used to detect and eliminate persons with a history of BCG vaccination, and described BCG policy and actual coverage in the population age group surveyed.

Data assembled from selected surveys:

The area population represented and the survey period.

In the youngest age ranges for which sufficient numbers were tested and read, the number tested and read, the number counted as infected with M. tuberculosis, and the mean age.

Information documenting quality of the data.

Criterion of infection

Induration reaction size distributions observed in each survey were used together with general knowledge about the specific and nonspecific components of observable distributions to set a uniform criterion for counting infected persons in order to improve comparability between surveys.

Because it can be assumed that nonspecific reactions to intermediate strength purified protein derivative tuberculins are only infrequently larger than the mode of the distribution of true reactions, an observed distribution of tuberculin reactions will be nearly free of contamination by nonspecific reactions above its mode.

Because of this, and because distributions among truly infected persons are symmetric, the number truly infected in a distribution contaminated by nonspecific reactions would be closely approximated by adding the number at the mode plus twice the number with reactions larger than the mode.

This method, described by Nyboe (1960) and by Comstock et al. (1971), was used to determine the number of infected persons in the selected groups when detailed reaction size distributions were available and when the mode of reactions among infected persons in the same general population could be estimated from distributions of either bacteriologically confirmed cases or older age groups in the population.

When such data were not available, a cutoff criterion was accepted as the second choice if it appeared reasonably adequate based on distribution data.

Prevalence

Prevalence at a particular calendar time for an age group was calculated as the number infected divided by the number tested and read. Reported prevalence figures were used when numerator or denominator counts were not specified.

Risk of infection

For comparability, a single uniform method was chosen to derive the approximate average annual risk of infection (Styblo et al, 1969, pg. 14). By this method, the annual risk of infection (R) for a group of average age (A) was derived from the prevalence (P) by

$$R = 1 - (1 - P)^{1/A}$$

The slope per year (B) of the trend between two risk estimates R_i and R_j at years Y_i and Y_j was approximated by

$$B = 1 - (R_j/R_i)^{1/T} \text{ where } T = Y_j - Y_i.$$

Average age

Average age was calculated by taking the midpoint of a single year (e.g., 6.5 for age given as 6 at last birthday) or a range of years (e.g., 7.0 for ages given as 6 to 7, or 7.5 for ages given as 5 to 9). When available, the weighted average age was calculated from year-of-age specific frequencies for an age group.

Time at which average risk occurred

For purposes of calculating and plotting trends, the calendar time of occurrence of a particular average annual risk (Y_R) was considered to center approximately at the calendar time of the midpoint of the average lives of the individuals in the age-group. This was calculated by $Y_R = Y_S - A/2$, where A is the average age of the group and Y_S is the midpoint time of the survey. For example, a group of average age 6.1 years surveyed at 1978.9 was taken to contribute information about the annual risk of infection for a time interval centered at $1978.9 - 6.1/2 = 1975.85$.

The midpoint time of the survey (Y_S) was calculated from the month, if stated, or the calendar year of the survey. For example, a survey stated to have taken place in 1975 was counted as occurring at 1975.5. A survey stated to have taken place from 1981 to 1982 was counted as occurring at 1982.0. A survey stated to have taken place from May 1952 to September 1953 was counted as occurring at 1953.04 (the midpoint between $4.5/12 + 1952$ and $8.5/12 + 1953$).

III. RESULTS

A variety of tuberculin skin-test surveys were found in the published and unpublished literature: National sample surveys, sample surveys of other large populations, school surveys, school BCG campaigns, and mass BCG campaigns.

Twenty-five countries were judged for the purposes of this project to have been adequately surveyed in whole or significant part since 1975.

Figures 1 and 2 display the results for each country by region. Table 1 contains the data used in the figures and describes the area population represented, the survey design, the level of BCG coverage in the age-group surveyed, and the antigen and criterion defining infection.

An arithmetic scale is used in Figure 1 in order to display the relative magnitudes of current estimates of annual risk of infection.

A logarithmic scale of annual risks is used in Figure 2 in order to represent constant percentage decreases or increases between current estimates and prior estimates as straight line trends.

Dark symbols in the figures indicate risk estimates based on 42 well-conducted national sample surveys in 16 countries. In Figure 2, dark solid lines are used to connect rates which are based on more than one large sample survey of the same country.

Light symbols in Figure 2 indicate risk estimates based on data representative of lesser populations.

Dark dashed lines in Figure 2 are used to connect risk estimates based on national sample survey data with risk estimates based on other substantial data such as tuberculin testing in the mass BCG campaigns. A light solid line was used to connect risk estimates from repeated surveys representative of lesser populations. A light dashed line was used to connect surveys of approximately the same population done in different years.

The symbol + was used in Figure 2 to indicate the average of a set of risk estimates which together would represent a larger population. Stratum estimates within sample surveys, plotted as separate points to reveal variation in risk, were displayed as a single estimate for the whole population in this way. Sets of small area estimates that were less representative of the whole population than would have been provided by formal probability sampling, such as the set of completed sampling units in a not yet completed national survey, or areas selected by judgement from geographic or economic strata in order to be representative of a large heterogeneous population, were also combined in this way.

To indicate the extent of underlying variability, risk estimates from single surveys that were representative of only small areas are indicated in Figure 2 by unconnected light symbols.

Results by region

Africa. The most recent data for the eight countries included indicate current annual risks of infection around 1 to 2 percent. In the five countries for which comparable earlier surveys were available, trends over 20 to 25 years appear to be downward at 1 to 6 percent a year.

Data for the nearly completed national survey of Tanzania indicate an annual risk of infection of 1.1 percent, based on a population weighted average pooling of the samples. However, comparisons with earlier data available for three of the 18 regions in the sample do not together suggest a downward trend.

Surveys in Botswana and Lesotho suggest downward trends of about 6 and 1 percent a year, respectively. In the most recent survey in Botswana, however, BCG coverage in the age group surveyed had reached 83 percent, leaving only 17 percent who had by whatever circumstance missed vaccination, upon which to base the estimate.

No national surveys were available for Cameroon, but two surveys of the capital city, Yaounde, were available. These surveys indicate a level of annual risk of 0.6 percent in 1980 after a 20-year declining trend of 3.6 percent a year.

Survey data collected 30 years apart were available for Addis Ababa, the capital city of Ethiopia. These surveys indicate a 1.3 percent level of annual risk in 1979 succeeding a downward trend of 3.7 percent a year from the 4.1 percent risk level in 1949. A survey of selected villages in the Southwest, however, indicated a 3.8 percent annual infection risk in 1973. This suggests that within Ethiopia there is large variability in the level and perhaps also in the trend of infection risk.

Data from the 1976 sample survey of Gambia indicates an annual risk in 1971 of 1.9 percent. The trend of risk could not be assessed in Gambia, however. The only prior data available are from the 1958-59 survey of the capital city, Bathurst. The annual risk of infection in the urban stratum (4.4 percent) was higher than in the rural stratum (1.7 percent), which was nearly the same as the national level in 1971.

The annual risk of infection derived from skin test data for all military recruits in Burundi in 1981-82 was 1.2 percent. Data from an earlier survey among a scattered provincial population were selected in order to provide a rough comparison. The annual risk of infection in that group was 2.7 percent in 1955.

No comprehensive survey of the Nigerian population was available, but data from a survey of one emirate indicates an annual risk of 2.3 percent in 1973.

Americas. National sample survey data were available only from Argentina, but repeated surveys of large local populations within Argentina and also within Brazil were available. In the latest surveys, infection risks in these local populations varied ten-fold from 0.2 to 2 percent per year. Over time in these local populations, there was evidence of a steep decline in risk in Argentina, and a lesser decline in Brazil.

The annual risk of infection derived from the 1974-78 survey of the entire population of Argentina was 0.6 percent. Data from three surveys repeated in Santa Fe City and three surveys repeated in areas around Santa Fe City indicate low recent levels in that region of 0.2 to 0.3 percent annual risk, and indicate that the trend of the previous 12 to 15 years was steeply downward at a rate of 7 to 10 percent a year. The two trends described by the two sets of three survey points suggest that the decline was much steeper between the first two surveys but that the trend was nearly level during the 5 years between the second and third surveys. In the 1960-61 sample survey of another area of Argentina, Resistencia Province, the annual risk of infection was 0.5 percent.

Capitals of Brazilian States and Territories were surveyed in 1970-73 and again in 1980. The annual risk of infection was 0.8 to 1.9 percent in 1976 with evidence of a yearly downward trend of around 2 to 3 percent. A 1983 survey of the provincial population of Rio Grande Do Sol indicated a 0.4 percent annual risk of infection, with a 2.4 fold higher risk in southern than in northern rural areas.

South-East Asia. Only Thailand had national survey data available, but repeated surveys of local area populations were available for both India and Indonesia. In these populations, annual risks in the decade of the 1970's appear to have ranged from about 1 to 5 percent. Annual risks were highest in Indonesia.

Based on the 1977-79 national survey, the annual risk of infection in Thailand was 2.2 percent in 1974. Two earlier surveys, one of Bangkok and Chiengmai Province in 1960-64, and one of localities of Chiengmai and Kanchanaburi Provinces in 1954, indicate annual risks only slightly higher than in the national survey 16 to 24 years later.

In India, three large local populations were twice surveyed. Declining trends of 1.6 and 3.5 percent a year were apparent in two of these populations, but an increase of about 1 percent was apparent in the third population. Village surveys in 7 regions throughout India in 1972 indicate annual risks on the order of 1 to 2 percent. The village survey in Kashmir, compared to a 1978 sample survey of the Kashmir Valley, suggest a declining annual trend, but the populations may be only nominally comparable.

Indonesia has followed a policy since 1974 of withholding BCG and conducting 5-yearly surveys in nine special areas selected to be situated throughout the country. Three surveys have been completed in five of these areas, two in three areas, and one in another area. There are also three surveys available from an area with low BCG coverage. The latest surveys in these areas indicate a median annual risk of 2.3 percent (range: 0.7 to 3.9). Earlier data from the 1964-65 survey in rural East Java indicate an annual risk of 1.6 percent. Comparing the earliest to the latest survey in each of the nine areas with repeated surveys, the median downward trend in the annual risk of infection was 2.5 percent a year (ranging from a decline of 6.3 percent to an apparent increase of 3.4 percent a year).

Eastern Mediterranean (and Algeria). For the purposes of this project, Algeria is grouped in this region instead of with the African countries. National survey data were available for six countries, and repeated surveys of local populations were available for a seventh country in this region. Recent annual risk levels appear to be low to very low in Bahrain, Libya, Algeria and Kuwait (0.2 to 0.5 percent), but high in Afghanistan and Pakistan (1.8 and 3.5 percent, respectively). Recent levels of risk in the Syrian cities of Aleppo and Homs also appear to be very low (0.3 and 0.14, respectively). The trend of risk also appears to have decreased steeply in the current low risk countries, but not as steeply in Pakistan and appears to have increased in Afghanistan.

Four local areas sampled in 1980-84 for the nearly completed national survey of Algeria had also been canvassed in the 1949-52 mass BCG campaign. Repeated surveys were also available for the population of Blida. Based on these comparisons, annual risk of infection appears to have decreased about 6 to 7 percent a year in Algeria. A similar trend is apparent between risk estimates for the Syrian cities of Aleppo and Homs.

In Kuwait between 1972 and 1981, all school entrants (including non-citizens) were tuberculin tested prior to BCG vaccination. Compared to the 1962-63 national sample survey estimate, the annual risk of infection appears to have declined by 10 percent a year. In Bahrain, a decline of nearly 12 percent a year appears to have occurred between the 1969 national sample survey and mass tuberculin testing of students in 1981. A similarly steep rate of decline may have occurred in Eastern Libya, if the areas and populations covered in the surveys are comparable.

The annual risk estimate based on the 1974-78 national sample survey of Pakistan, which was completed except for Baluchistan, compared to the estimate based on the mass BCG campaign of 1949-54, suggests that the risk of infection declined by 4.2 percent a year, but that the risk was still high (1.8 percent) in 1972. The risk estimate based on the partially completed 1961-62 national survey is consistent with this trend.

The annual risk of infection in Afghanistan appears high (3.5 percent in 1978), without having much changed since 1958, compared with an estimate based on tuberculin testing program data collected in 1963 in selected schools around the country.

Western Pacific. Eleven national sample surveys were available for five countries in this region. In the Republic of Korea, the annual risk of infection appears to be declining, but remains high at about 2 percent. The risk appears to be about 1.8 percent in the Philippines, based on the recent national survey. Infection risk appears to be relatively low in Malaysia (0.4 percent) and Samoa (0.5 percent).

In China the annual risk of infection appears to have been 1 percent in 1975, based on areas without BCG coverage in the very large 1979 national survey sample.

Based on infection risk estimates from the five 5-yearly national surveys available from the Republic of Korea, the trend appears to be downward at 3.8 percent a year to a level in 1984 of 2 percent.

IV. DISCUSSION

Interpretation of this collection of risk and risk trend estimates must take into consideration the many potential sources of error, both in determining absolute levels of risk and in comparing risk estimates among countries or between periods within the same country. Some of these sources of error appear to have been adequately controlled, but the influence of other sources of error (identified and unidentified) must be acknowledged as unknown.

Tuberculin skin testing is a usefully accurate technique for separating groups at relatively high and low risk of future Mycobacterium tuberculosis disease, but is liable to varying degrees of error in identifying all the infected and specifying all the uninfected.

Sensitization to environmental mycobacteria is a source of nonspecific reactivity to tuberculin that is especially prevalent in tropical regions (Edwards and Edwards, 1960). There was evidence in reaction size distributions of a high prevalence of nonspecific sensitivity in several of the African and Asian populations surveyed, but distribution data for older groups or for bacteriologically confirmed cases were frequently available for locating the mode of the distribution of true infections and deriving their number by assuming the distribution to be symmetric.

There is underlying variability in tuberculin reactivity as evidenced by the variation between simultaneously applied identical tests in the same individual (Chaparas et al, 1985). Although this reduces precision of estimation, it is not expected to introduce systematic error.

A variety of antigen preparations and dosages were used, and these differed between countries and within countries over time. Estimating the number infected by using the symmetry of the distribution of true reactions about the mode would adjust for differences in specific biological potency between antigen preparations, but would not be effective in controlling for large differences in specificity of antigens.

The Mantoux method of administration was virtually always used, and all the selected studies measured induration of reaction. Techniques of measurement varied somewhat, but almost always consisted of transverse measurement at 48 or 72 hours after administration.

Waning of tuberculin reaction size over time, and complete reversion to zero reactivity are recognized to occur (Sutherland, 1971). To the extent reversions have occurred in a population, tuberculin test prevalence at a point in time will underestimate the proportion who were ever infected. But limiting to groups of about the same low average age may have minimized and kept comparable the effects of cumulative reversion.

Boosting is a potential problem in survey designs that retest the same individuals (Styblo, 1984). Boosting is the effect of an initial tuberculin test which, although it elicits slight or no reactivity itself, apparently acts as an antigenic stimulus able to recall waned or reverted preexisting mycobacterial sensitivity so that a subsequent test does elicit reactivity that is indistinguishable from conversion caused by M. tuberculosis transmission in the interval between the tests. Although some of the data compiled in this project came from surveys using a repeated testing design, the problem of boosting was avoided by selecting age groups too young to have been tested in earlier rounds.

BCG vaccination, a major component of tuberculosis control activities in developing countries, compromises the usefulness of tuberculin testing as a means to survey risk of infection. Tuberculin skin testing cannot distinguish between sensitivity caused by BCG vaccination and sensitivity caused by natural infection with M. tuberculosis, because BCG vaccination, unlike nonspecific sensitization, causes reactions that are distributed with nearly the same mode and shape as reactions caused by true infections with M. tuberculosis. For this reason the risk of infection in the whole population can only be inferred from risk measured in the unvaccinated.

In some of the countries for which survey data were selected, the policy was to vaccinate indiscriminantly without using tuberculin testing to select the as yet uninfected. In these countries, surveys limited to children without evidence of vaccination are subject to bias only to the extent that children selected for vaccination, either by choice or by local availability, are at a different risk of infection than the unvaccinated. This is a potentially severe source of bias if BCG coverage is very high.

In some of the surveys selected for this project, none of those surveyed had been vaccinated, either because BCG had not been introduced into that locale, or because BCG was withheld by policy until a later age. In other surveys, the unvaccinated were identified by physical examination to detect the characteristic scar. Levels of coverage by BCG were very low in some surveys, in which case representation of the whole population by the unvaccinated would be unaffected. In a few surveys, BCG coverage was so high that representation was demonstrably affected.

In other countries, the policy was to tuberculin test and vaccinate only those with weak or no reactions. In this situation, a later survey among children without BCG scars would be liable to severe overestimation bias. This was avoided by selecting age groups before the age of eligibility for selective vaccination programs in effect in a particular country.

Failure to test and measure every person in a survey sample can introduce a bias if those not reached by the survey are at a different risk of infection than those who were reached. A severe bias might also result from failure to measure a high proportion of those tested, because whether a survey participant returned for reading could well depend on whether there had been a reaction. Surveys with severe problems of this kind were not selected. Adequate coverage was a criterion for selection by this project, and was high in most of the surveys selected.

Although the average annual risk for a group lies in the calendar interval from birth to the time of the survey, it need not be at the interval midpoint. If the risk is changing substantially over time, as data from several of the surveys indicate, or varies with age, as has been suggested by the work of Sutherland (1976), and as might be expected in some of the diverse cultures surveyed, a complex model is necessary to estimate the risk at a particular time (Styblo, 1969). Limiting to young, narrow age groups was used as a way to bracket risk estimates within usefully narrow time intervals.

Average annual risk estimates for the same populations surveyed in different years are connected with straight lines in Figure 2 to indicate the average trend that would result in the observed risk estimates. An exponential trend is to be expected in the situation of a long-term constant ratio of the number of new cases produced by each existing case (Sutherland, 1976). However, without data for intermediate years, it is difficult to know the true shape of a trend line. In some instances where more than 2 surveys were available, the trend appeared to follow a fairly constant geometric decrease, but in other instances, the trend appeared to have changed over time.

Many survey reports presented more detailed analyses of infection risk and its trend, but for the purposes of this review, methods were adopted that could be simply and consistently applied to all data. In the few instances of large disagreement, the present analysis may be in error.

V. CONCLUSIONS

This attempt to review and compile data available from major surveys carried out in developing countries since 1975 falls far short of providing a comprehensive picture of tuberculous infection risk and trend in the entire developing world. The assembled data cannot safely be generalized beyond the specific target populations surveyed. However, the assembled data do provide several recent objective examples of apparent progress and lack of progress in tipping the balance against the tuberculosis epidemic.

The review found that since 1975 many large scale surveys have been successfully carried out in the developing countries, with impressive attention to correct procedure and technique.

In most of the countries repeatedly surveyed in whole or in part, risk of infection does appear to have diminished. In some of these countries, the decline appears to have been substantial, and current levels of risk have become very low in a few countries.

In other countries, however, rates of infection appear to remain high and there is no indication of decrease.

It may be speculated that general improvements in living standards and tuberculosis control programs are having an effect against the tuberculosis epidemic in some parts of the developing world as it already has in the more developed countries. In the rest of the developing world, however, very little is known of the current magnitude and trend of the risk of tuberculous infection.

REFERENCES

- Bleiker, M.A. & Styblo, K. (1978) The annual tuberculosis infection rate and its trend in developing countries. Bulletin of the International Union Against Tuberculosis, 53, 295-298.
- Comstock, G.W., Furculow, M.L., Greenburg, R.A., Grzybowski, S., Maclean, R.A., Baer, H. & Edwards, P.Q. (1971) The tuberculin skin test. American Review of Respiratory Disease, 104, 769-775.
- Chaparas, S.D., Vandiviere, H.M., Melvin, I., Koch, G. & Becker, C. (1985) Tuberculin test. Variability with the Mantoux procedure. American Review of Respiratory Disease, 132, 175-177.
- Edwards, P.Q. & Edwards, L.B. (1960) Story of the tuberculin test from an epidemiologic viewpoint. American Review of Respiratory Disease, 81, suppl., 1-47.
- Nyboe, J. (1960) The efficacy of the tuberculin test. Bulletin of the World Health Organization, 22, 5-37.
- Pio, A. Epidemiology of tuberculosis (1984) Minerva Medica, 75, 507-517.
- Styblo K., Meijer, J. & Sutherland, I. (1969) The transmission of tubercle bacilli. Its trend in a human population. Tuberculosis Surveillance Research Unit Report No. 1. Bulletin of the International Union Against Tuberculosis, 62, 1-104.
- Styblo K. (1984) Epidemiology of tuberculosis. Infektionskrankheiten und ihre Erreger, VEB Gustav Fischer Verlag, Jena.
- Styblo K. The relationship between the risk of tuberculous infection and the risk of developing infectious tuberculosis. Bulletin of the International Union Against Tuberculosis, 60, 117-119.
- Sutherland I. (1971) The effect of tuberculin reversion upon the estimate of the annual risk of infection. Bulletin of the International Union Against Tuberculosis, 45, 115-118.
- Sutherland I. (1976) Recent studies in the epidemiology of tuberculosis, based on the risk of being infected with tubercle bacilli. Advances in Tuberculosis Research, 19, 1-63.

References to Table 1 by Region and country

WHO African Region - Region Africaine de l'OMS

Algeria

(a) Evaluation of the risk of tuberculous infection in a country with a high BCG-coverage in newborns. Methodology of the national tuberculin survey in Algeria. R Amrane. ITSC 1985 Progress Report, Vol. 2.

(b) Method for evaluating the infection risk (and its trend) in a country with a high prevalence of tuberculosis and where generalised BCG is applied at birth to newly born infants. R. Amrane, H. Ait-Mesbah, P. Chaulet. Bulletin of the International Union Against Tuberculosis, 59, 141-143, 1984.

(c) Mass BCG vaccination in Algeria, 1949-52, with special reference to statistics on tuberculin testing and BCG vaccination. Jorgen Nyboe, Mette Soegaard. Tuberculosis Research Office, WHO, Copenhagen. Published by The International Tuberculosis Campaign, December 1953.

(d) Résultats des mesures de surveillance de la Tuberculose en Algérie de 1980 à 1985. R. Amrane, H. Ait Mesbah, M.T. Hani. Reported at XXVIème Conférence Mondiale de l' Union Internationale contre la Tuberculose. Singapore, 2-7 novembre 1986.

Botswana

(a) Prevalence of tuberculosis in Botswana and Lesotho: Results of two random sample surveys. P.B. Fourie & K. Knoetze. The South African Journal of Epidemiology and Infection, 1, 32-37 (1986).

(b) Tuberculosis in Botswana. Results of an epidemiological survey 1981. E.T. Maganu, Epidemiology Unit, Ministry of Health, P.O. Box 10004, Gabarone.

(c) Tuberculosis survey in Basutoland, Bechuanaland, and Swaziland. WHO Tuberculosis Research Office, Copenhagen, April 1958.

Burundi

(a) Chapitre 1. Ampleur du problème de la TBC au Burundi - calcul de l'incidence réelle. Service d'Intégration de la Lutte Contre la Lèpre et la Tuberculose. Lutte Nationale Contre la Tuberculose, Rapport Annuel 1984. Ministère de la Santé Publique, République du Burundi.

(b) A tuberculin sensitivity survey in Burundi. June - September 1964. WHO Regional Office for Africa, P.O. Box 6, Brazzaville.

Cameroon

(a) Intérêt des enquêtes tuberculiques par sondage. A propos d'une enquête réalisée dans les écoles de Yaoundé: Méthodologie et résultats préliminaires. H.G. Delolme, P. Blin, G. Roscigno, M. Merlin, G. le Mao, L. Sentilhes. XV Conf. Tech. OCEAC 1984, 28-34. Organisation de Coordination pour la Lutte Contre les Endémies en Afrique Centrale, Secrétariat Général, B.P. 288, Yaoundé, République du Cameroun.

(b) A tuberculin sensitivity survey in Cameroon. March - May 1964. WHO Regional Office for Africa, P.O. Box 6, Brazzaville

Ethiopia

- (a) ITSC 1983/84 Report & 9th Progress Report. M.A. Bleiker, N. Dow, H. Ypma, H. Meesters. ITSC.
- (b) A tuberculin skin test survey in Southwestern Ethiopia. G.K. Fuller, N. Gameda, D. Fuller, V. Demerest. Tropical and Geographical Medicine, 31, 365-373 (1979).
- (c) Control of tuberculosis in Ethiopia. P. Chasles, A. Octapodas. Ethiopian Medical Journal, 1, 128-133, (1963).

Gambia

- (a) Tuberculin survey in Gambia. ITSC 5th Report.
- (b) Skin sensitivity to human PPD and PPD-marianum in schoolchildren in the Gambia West-Africa, 1976.
- (c) Tuberculosis survey in Gambia. MHO/PA/23.60. WHO Regional Office for Africa, Brazzaville, February 1960.

Lesotho

- (a) Prevalence of tuberculosis in Botswana and Lesotho: Results of two random sample surveys. P.B. Fourie, K. Knoetze. The South African Journal of Epidemiology and Infection, 1, 32-37 (1986).
- (b) Tuberculosis survey in Basutoland, Bechuanaland, and Swaziland. WHO Tuberculosis Research Office, Copenhagen, April 1958.
- (c) Report on the initial examination in the WHO-assisted tuberculosis control project Basutoland-2. Report to the Regional Director, AFRO from Dr Anton Geser, SMO AFRO-53.

Nigeria

- (a) Skin sensitivity to human PPD and to PPD-marianum in schoolchildren in the Kazuare Emirate in the Kano State of Nigeria 1977. M.A. Bleiker, G.L. Pape, O. Misljenovic, & I. Blijker. International Tuberculosis Surveillance Center.

Tanzania

- (a) Tuberculin survey within the national tuberculosis/leprosy programme in Tanzania. ITSC 9th Progress Report, 1984.
- (b) Tuberculin survey in the Shinyanga Region, the United Republic of Tanzania, 1979. ITSC 7th Progress Report, 1980.
- (c) Draft report. Tuberculin survey and tuberculin training in the Morogoro- and Arusha-Regions in the United Republic of Tanzania. ITSC 1982/83 Progress Report.
- (d) The risk of tuberculosis infection in the Dodoma Region, Tanzania. Results of a tuberculin survey among schoolchildren carried out in November and December 1978 by a team of the TB/Leprosy Unit of the Ministry of Health and the Dodoma Regional Leprosy/TB Scheme. Report to Dr Antonio Pio, WHO TRI Unit. Dr Jaap F. Broekmans, 12 December 1978, P.O. Box 876, Dodoma.
- (e) The national tuberculin survey, Tanzania. M.A. Bleiker, S.J. Nkinda, O. Misljenovic, D.W. Mulder & K. Styblo. TSRU Davos Meeting, October 1985.

(f) Skin sensitivity of human PPD in schoolchildren in Tanga in the United Republic of Tanzania, 1977. M.A. Bleiker, Dr Madundo, O. Misljenovic, M.H. Millia, I.F. Abdallah. ITSC.

(g) Tuberculosis programme implementation: Tanzania. K. Styblo. TRI/TB:PHC/86.12
22-26 September 1986.

(h) The National Tuberculosis/Leprosy Programme in Tanzania. Summary Report no. 17 on the visit to Tanzania, February 1987. K. Styblo.

(i) Tanzania national tuberculin survey. M.A. Bleiker, H.J. Chum, S.J. Nkinda, K. Styblo. TSRU/IUAT Progress report 1987, Volume 2, 135-146.

Region of the Americas - Région des Amériques de l'OMS

Argentina

(a) Determination del Riesgo de Infección Tuberculosa en la Republica Argentina. Resultados de la primera encuesta tuberculínica 1974 - 1978. Ministerio de Bienestar Social, Secretaría de Salud Pública. Instituto Nacional de Tuberculosis, Recreo, Santa Fé, Argentina

(b) The tuberculosis situation in Argentina. PAHO Epidemiological Bulletin, Vol. 4, No. 3, 1983, 8-10.

(c) Official Communication from Dr Eduardo Balestrino, Director Instituto Nacional Epidemiologica, 10 de marzo de 1986.

(d) Informe sobre la encuesta de prevalencia de la tuberculosis en la Provincia del Chaco. Ministerio de Asistencia Social y Salud Pública de la Nación. Organización Mundial de la Salud, UNICEF, Neuquén, Marzo 1961.

Brazil

(a) Estudo da prevalência de infecção tuberculosa na população escolar de 6 - 7 anos no Rio Grande do Sul. Resultados do 1º tempo, 1983 -1988, Volume 1. Estado do Rio Grande do Sul, Secretaria da Saude e do Meio Ambiente, Porto Alegre, 1984.

(b) Informe final sobre la consultoria en analisis de informacion epidemiologica y operational del programa nacional de control de tuberculosis, Division Nacional de Pneumologia Sanitaria Ministerio de Salud, Rio de Janeiro, Brasil
Del 8 de noviembre al 6 de diciembre de 1981. Dr E. Balestrino, Consultor, Organización Panamericana de la Salud.

WHO South-East Asia Region - Region de l'Asie du Sud-Est de l'OMS

India

(a) Assignment report on a tuberculosis longitudinal survey, National Tuberculosis Institute, Bangalore, WHO Project: India O103. Dr T. Olakowski, Senior Medical Officer (in cooperation with the members of the Technical Co-ordination Committee at the National Tuberculosis Institute, Bangalore), January 1969 - March 1972. SEA/TB/129, 12 September 1973 Restricted. SEA-73/2584.

(b) Tuberculosis in a rural population of South India: Report on five surveys. A.K. Charaborty, H. Singh, K. Srikantan, K.R. Rangaswamy, M.S. Krishnamurthy, J.A. Stephen. Ind. J. Tub., Vol. XXIX, No.3, 1982, 153-167.

(c) Some aspects of a tuberculosis prevalence survey in a South Indian district. Raj. Narain, A. Geser, M.V. Jambunathan, M. Subramanian. Bull WHO, 1963, 29, 641-664.

- (d) Prevalence of infection among unvaccinated children for tuberculosis surveillance. A.K. Chakraborty, K.T. Ganapathy, G.D. Gothi. Indian J Med Res, 72, July 1980, 7-12.
- (e) Changes in the prevalence rates of infection in younger age groups in a rural population of Bangalore District over a period of 5 years. A.G. Kurthkoti, Hardan Singh. National Tuberculosis Institute Newsletter (1985) 21/2, 28-40.
- (f) Prevalence of tuberculosis in a South Indian District - twelve years after initial survey. G.D. Gothi, A.K. Chakraborty, S.S. Nair, K.T. Ganapathy, G.C. Banerjee. Ind. J. Tub., Vol. XXVI, No. 3, 122-135.
- (g) Tuberculosis prevalence survey in Kashmir valley. S. Mayurnath, D.S. Anantharaman, G.V.J. Baily, M.P. Radhamani, R.S. Vallishayee, P. Venkataraman, S.P. Tripathy. Indian J Med Res 80, August 1984, 129-140.
- (h) Prevalence of non-specific sensitivity in some parts of India. Raj Narain, M.S. Krishnamurthy, D.S. Anantharaman. Indian J Med Res, 63, 8, August 1975, 1098-1109.
- (i) Tuberculosis prevalence survey in Tumkur District. Raj Narain, A. Geser, M.V. Jambunathan, M. Subramanian. Ind. J. Tub., Vol. X, No. 3, 87-116.

Indonesia

- (a) The risk of tuberculosis infection in the District of Tangerang (Indonesia) derived from the results of tuberculin testing of schoolchildren aged 7 to 10 years, 1972-1983. ITSC 1983/84 Report.
- (b) Tuberculin resurveys in two areas in Indonesia. ITSC 1981/82 Report.
- (c) Tuberculosis Research and Control - Control of bacterial and intestinal diseases. Assignment Report, 1 December 1981 - 28 February 1982. Dr Yoshikumi Azuma, WHO Short-Term Consultant. WHO Project INO BVM 001. SEA/TB/167, 30 September 1982, Restricted.
- (d) Tuberculosis control in Indonesia 1952-65. Report on WHO projects: SEARO 0003 and Indonesia 0050. Section I: Text, Section II: Data Tables. Prepared by WHO Regional Office SEA/TB/92 I,II, SEA/VHS/95 I,II, 19 December 1968. Restricted SEA-68/2638.
- (e) Annual tuberculosis infection rates between 1972 and 1978 in the district of Tangerang (Indonesia) based on the results of tuberculin skin testing of schoolchildren aged 7-10 years. ITSC 6th Progress Report.
- (f) Skin sensitivity to human PPD and to avian PPD in schoolchildren in Indonesia, 1972. H. Kusnadi, A.S. Gunardi, M.A. Bleiker. ITSC Confidential Report in WHO files.
- (g) 1986 Report of Dr Tripathy.
- (h) ITSC working papers.

Thailand

- (a) Epidemiology of tuberculosis in Thailand. Suchart Daramas, MD. Report in the WHO TRI Unit files.
- (b) Situation of tuberculosis in Thailand. Results of first national epidemiological survey conducted in Bangkok and the Province of Chiangmai during 1960 - 1964. Proceedings of Eastern Regional Meeting, Bangkok 1964.
- (c) Data for the assessment of naturally acquired tuberculin sensitivity in seven countries of Asia. WHO Tuberculosis Research Office, Copenhagen, June 1955.

(d) Tuberculosis epidemiology in Thailand. Dr Boonsong Sunakorn, J Med Assoc Thailand, 32, 1969, 157-162.

WHO Eastern Mediterranean Region - Region de la Mediterranee Orientale de l'OMS

Afghanistan

(a) EM/TB/159 EM/AFG/BVM/001 February 1983. Assignment Report. National Tuberculosis Survey in Afghanistan, 3 August - 2 November 1982, by Dr K.S. Aneja, WHO Consultant.

(b) SEA/TB/96, 10 September 1969. Restricted. Assignment Report on Tuberculosis Advisory Services (WHO Project: Afghanistan 0033 (UNDP/TA) January 1965 - July 1968 by Dr G. Khan, WHO Medical Officer.

(c) EM/SEM.TB/4.1, 20 August 1975. Regional Seminar on Recent Trends in Tuberculosis Control, Karachi, 23-30 October 1975. Tuberculosis Control in the Eastern Mediterranean Region. Review of the Present Situation by Dr J. Kaleta, WHO Regional Adviser on Tuberculosis.

Bahrain

(a) Survey of tuberculous infection in schoolchildren in Bahrain. M. Ilyas Khan, Tubercle, 63 (1982), 287-289.

Kuwait

(a) Kuwait National Tuberculosis Control Program. Dr Mohammed Abdel Aty, Head, Tuberculosis Control Unit. The 18th Middle-East Regional Conference of the IUAT, 5-8 February 1983, Kuwait.

Libya Arab Jamahiriyah

(a) Assignment Report. A national tuberculosis prevalence survey in the Socialist People's Libyan Arab Jamahiriyah, February 1976 - December 1977. Dr Syed Ali Husain, WHO Medical Officer and Mr Ib Thorup, WHO Statistician. EM/ST/121, EM/TB/152, EM/LIY/BVD/001, August 1978.

(b) Regional tuberculosis prevalence survey. Report on Cyrenaica, Libya, 11 July to 22 September 1959. Epidemiological and Statistical Centre, WHO Regional Office for the Eastern Mediterranean, Alexandria, UAR. EM/TB/58, EM/ST/14, February 1961.

(c) Report on BCG assessment work in Libya. WHO Tuberculosis Research Office, Copenhagen, April 1956.

Pakistan

(a) Epidemiological situation of tuberculosis in Pakistan. Results of the National Tuberculosis Prevalence Survey 1974-1978. Dr Jan Kaleta, Mr Nisar, A. Chaudhry. TB Seminar, Lahore, April 1982.

(b) Travel Report by Dr Antonio Pio, CDS/TRI, 21.1.85 T9/370/12PAK.

(c) Report of the tuberculosis survey in Karachi, Rawalpindi & Lahore Division of West Pakistan. Directorate of Tuberculosis Control (Health Division), Government of Pakistan, October 1962.

(d) BCG-vaccination programme in Pakistan. E. Roelsgaard, H. Christensen, E. Iversen. Bull. Wld Hlth Org., 1957, 17, 187-202.

(e) Assignment Report on a visit to Pakistan Tuberculosis Control Programme. 9 May - 6 June 1983.

Dr Jan Kaleta, WHO Consultant. EM/TB/161, August 1983.

Syrian Arab Republic

(a) The annual risk of tuberculous infection in the Syrian Arab Republic, 1950-1983. ITSC 1983/84 Report.

(b) Estimates of the risk of infection in schoolchildren in Syria between 1938 and 1978. ITSC 7th Progress Report 1980.

(c) Tuberculosis survey in the Syrian Arab Republic. Epidemiological and Statistical Centre, WHO Regional Office for the Eastern Mediterranean, Alexandria, UAR, April 1962.

(d) Mass BCG vaccination in Syria, 1950, with special reference to statistics on BCG vaccination and pre- and post-vaccination allergy. Published by The International Tuberculosis Campaign, care of UNICEF, 24 Rue Borghèse, Neuilly-sur-Seine, France.

(e) Skin sensitivity to human PPD in schoolchildren in the Governates of Aleppo, Homs, and Raqua in the Syrian Arab Republic, 1983. M.A. Bleiker, O. Misljenovic, O. Post, I. El-Rifai, B. El-Kateb, W. Sankari, K. Nahawi, N. Nasser, F. Kana, B. Hanifi, S. Trabalsi, R. Assaf. ITSC Report.

WHO Western Pacific Region - Region du Pacifique Occidental de l'OMS

China

(a) Nationwide random survey for pulmonary tuberculosis conducted in 1979. The National Tuberculosis Control and Research Centre, September 1985. Draft for possible publication, submitted to TRI Unit, WHO.

(b) Nation-wide survey for the epidemiology of pulmonary tuberculosis. Conducted in 1979. Zhonghua Jiehe He Huxixi Jibing Zazhi, 1982, 5, 67-70 (in Chinese).

Malaysia

(a) An analysis on the first community tuberculin survey in peninsular Malaysia (1976-1977). Dr H.T. Lin and Mr Ib Thorup. National Tuberculosis Programme of Malaysia. ICP/BVD/001-E(Malaysia), 18 February 1979.

Philippines

(a) Report on a national tuberculosis prevalence survey in the Republic of the Philippines 1981-1983. National Institute of Tuberculosis, June 1984.

(b) Second final report from the Philippines BCG mass vaccination programme. Dr Anton Geser, BCG Consultant. Manila, 1 August 1955. Report in the WHO TRI Unit files.

Republic of Korea

(a) Report on the 5th tuberculosis prevalence survey in Korea, 1985. Ministry of Health and Social Affairs, Korean National Tuberculosis Association.

(b) Assignment report. Tuberculosis control, Republic of Korea, 5 February to 3 March 1986. Dr Toru Mori, WHO Consultant (WP)CHD/ICP/TUB/001 E, 3 November 1986.

(c) The value of periodical tuberculosis prevalence surveys to assess the epidemiological trend of the problem in developing countries. Ingela Sjögren. ITSC 1984 Progress Report, Vol. 1.

(d) Report on the 4th Tuberculosis Prevalence Survey in Korea. 1980. Ministry of Health and Social Affairs, Korean National Tuberculosis Association.

(e) Prevalence Survey in Korea, B.W. Jin, The Korean National Tuberculosis Association, Korean Institute of Tuberculosis, 1984. Report in the WHO TRI Unit files.

(f) Summary report on a national tuberculosis prevalence survey in Korea. (June - November 1965). WHO Regional Tuberculosis Advisory Team, WHO Regional Office for the Western Pacific, Manila, Philippines. WHO/TB/Techn.Information/67.57.


Samoa

(a) Report on the tuberculosis/leprosy prevalence survey in Samoa, 26 June - 24 October 1975. WHO Regional Tuberculosis Control Team and WHO Leprologist. ICP/BVD/001-E (Western Samoa) 5 December 1977.

(b) Assignment report. Tuberculosis control, Western Samoa, 18 September to 7 July 1974. Dr Nak Chin Chung, WHO Consultant Western Samoa. 1201-E (WES/MBD/01), 25 October 1974.


(c) Assignment report. Tuberculosis control, Western Samoa, 8 March 1980 to 20 December 1981. Dr Qian Yuan Fu, WHO Medical Officer. SMA/BVM/002-E, 13 April 1982.

RISK (%)





LEGEND:

- ◇ AFRICA
- AMERICAS
- ⊕ SOUTH-EAST ASIA
- △ EASTERN MEDITERRANEAN
- WESTERN PACIFIC





RISK (%)





RISK (%)





LEGEND

▽ ○ ⊕ × □ △ ◇ ✱ Survey estimates of risk

▽ ○ ⊕ × □ △ ◇ ✱

+ + Average of risk estimates

———— Trend between risk estimated
- - - - - in different years

————
.....

Dark symbols and lines are used to display risk estimates and trends for national populations and large parts of national populations.

Light symbols and lines are used to display risk estimates for subnational units.

Solid lines connect surveys that closely represent the same population.

Dashed lines connect surveys that represent approximately the same population.

TABLE 1. ANNUAL RISK OF TUBERCULOUS INFECTION IN DEVELOPING COUNTRIES
BASED ON TUBERCULIN SURVEYS

I. WHO African Region (except Algeria) - Region Africaine de l'OMS (sans Algerie)

Year	Area Represented	Survey design	BCG (%)	Antigen and Infection Criterion	Survey Year Midpt.	Age (midpt.)	Tested N	Infect- ted N	Preva- lence (%)	Risk/ year (%)
<u>Botswana</u>										
1981-82 ^{ab}	Whole country	Sample survey	83.3	RT23 2TU, Mode=13	1982.0	0-14 (6.5)	257	...	8.17	1.30
1956-57 ^c	Whole country	Sample survey	0	RT19-20-21 5 TU, Mode=16	1957.0	0-14 (6.6)	1450	472	32.55	5.79
<u>Burundi</u>										
1982-84 ^a	Military recruits	All recruits	71.0	RT23 2TU, 9mm+	1983.5	18-22 (20.5)	912	195	21.38	1.17
1964 ^b	Northern section of Muramvya Province	Sample survey	56.4	RT23 2TU, 9mm+	1964.6	15-19 (17.5)	202	76	37.62	2.66
<u>Cameroon - Cameroun</u>										
1984 ^a	Yaounde	Sample survey of schools	58.4	PPD 71 10U, 9mm+	1984.5	7- 9 (8.5)	860	46	5.35	0.64
1964 ^b	Urban Yaounde	Sample survey	...	RT23 2TU, 9mm+	1964.3	5- 9 (7.5)	326	31	9.51	1.32
<u>Ethiopia - Ethiopie</u>										
1983 ^a	Addis Ababa	Sample of schools	49.4	RT23 2TU, 10mm+	1983.9	6-15 (8.6)	1251	133	10.63	1.30
1977 ^b	Southwest area	Selected villages 2TU, 10mm+	1977.5	6-10 (8.5)	185	52	28.11	3.81
1953-55 ^c	Addis Ababa	BCG school campaign	0	...	1954.7	7-14 (11)	37	4.1
<u>Gambia</u>										
1976 ^{ab}	Whole country	Sample survey	0.7	RT23 2TU, Mode=15	1976.4	6-16 (9.4)	2397	399	16.65	1.92
1958-59 ^c	Bathurst in 2 strata	Sample survey:	...	RT23 1TU, Mode=17	1959.1	5-14 (9.4)	16.38	1.89
	Urban stratum	at 50 per 1000					176	61	34.66	4.43
	Rural stratum	at 7 per 1000					400	61	15.25	1.74

I. WHO African Region (except Algeria) - Region Africaine de l'OMS (sans Algerie) (continued)

Year	Area Represented	Survey design	BCG (%)	Antigen and Infection Criterion	Survey Year Midpt.	Age (midpt.)	Tested N	Infect N	Prevalence (%)	Risk/year (%)
<u>Lesotho</u>										
1981-82 ^a	Whole country	Sample survey	70.4	RT23 2TU, Mode=14	1982.0	0-9 (5.2)	158	...	10.13	2.03
1956-57 ^b	Whole country	Sample survey	0	RT19-20-21 5TU, Mode=17,18	1957.0	0-9 (4.9)	1101	142	12.90	2.78
1962-65 ^c	Rural area around Maseru	Sample survey	0	RT23 1TU, Mode=19	1963.8	0-9 (4.9)	10216	1342	13.14	2.83
<u>Nigeria</u>										
1977 ^a	Kazaure Emirate of Kano State	School survey	32.1	RT23 2TU, Mode=15	1977.4	4-13*(7*)	15.0	2.29
<u>United Republic of Tanzania - Republique-Unie de Tanzanie</u>										
1983 ^{acehi j}	Arusha Region	Sample survey	52.4	RT23 2TU, Mode=17	1983.6	7-14 (10.0)	3125	307	9.82	1.03
1984	Coast (Pwani) Region	of schools	37.6		1984.8	(9.6)	681	74	10.87	1.19
1984	Dar-es-Salam Region	in progress,	45.0		1984.8	(9.7)	1338	137	10.24	1.11
1986	Iringa Region	90% completed	57		1986.4	(10.1)	1723	156	9.05	0.94
1986	Dodoma Region		49		1986.9	(10.7)	1743	358	20.54	2.13
1987	Kigoma Region		49.1		1987.7	(10.2)	1188	143	12.04	1.25
1985	Kilimanjaro Region		52.4		1985.4	(10.2)	2324	111	4.78	0.48
1984	Lindi Region		31.3		1984.6	(10.3)	2665	240	9.01	0.91
1986	Mbeya Region		64		1986.5	(10.9)	1621	193	11.91	1.16
1986	Morogoro Region		67		1986.6	(11.0)	865	158	18.27	1.82
1984	Mtwara Region		41.8		1984.5	(9.7)	629	122	19.40	2.20
1985	Mwanza Region		...		1985.8	(10.2)	2420	...	9.5	0.97
1986	Rukwa Region		67		1986.4	(10.8)	385	47	12.21	1.20
1984	Ruvuma Region		46.1		1984.5	(10.4)	1412	140	9.92	1.00
1985	Shinyanga Region		...		1985.4	(9.8)	3666	...	11.6	1.25
1985	Singida Region		...		1985.7	(10.5)	985	...	16.3	1.68
1987	Tabora Region		...		1987.7	(11.2)	1794	...	8.3	0.77
1983	Tanga Region		50.1		1983.8	(10.4)	2462	238	9.67	0.97
1983-87	(18 Area weighted average)				1985.6	(10.3)	30982	...	10.9	1.11
1979 ^b	Shinyanga Region, except Shinyanga Distr.	Sample survey	50.9	RT23 2TU, 10mm+	1979.5	6-14 (10.0)	1817	356	19.59	2.16
1978 ^d	Dodoma Region	Sample survey	63.3	RT23 2TU, Mode=17	1978.9	6-14 (10.2)	1329	166	12.49	1.30
1977 ^e	Tanga Town and nearby rural area	Sample survey of schools	79.0	RT23 2TU, Mode=17	1977.1	7-14 (9.7)	383	33	8.62	0.92

II. WHO Region of the Americas - Region des Ameriques de l'OMS

<u>Year</u>	<u>Area Represented</u>	<u>Survey design</u>	<u>BCG (%)</u>	<u>Antigen and Infection Criterion</u>	<u>Survey Year Midpt.</u>	<u>Age (midpt.)</u>	<u>Tested N</u>	<u>Infect- ted N</u>	<u>Preva- lence (%)</u>	<u>Risk/ year (%)</u>
<u>Argentina - Argentine</u>										
1974-78 ^{ab}	Whole country	Sample survey	...	RT23 2TU, 10mm+	1976.5	6- 7 (6.5)	26902	970	3.61	0.56
1983 ^c	Areas around	Sample survey	81.6	..., 10mm+	1983.5	5- 8 (7.0)	325	...	1.8	0.26
1979-80 ^c	Santa Fe City		63.7	..., 10mm+	1980.0	6- 7 (7.0)	443	...	2.0	0.29
1967-68 ^c			31.6	..., 10mm+	1968.0	5- 8 (7.0)	1221	...	8.8	1.31
1979-80 ^c	Santa Fe City	All schools	57.7	..., 10mm+	1980.0	6- 7 (7.0)	2125	...	1.6	0.23
1974-75 ^c			12.9	..., 10mm+	1975.0	6- 7 (7.0)	3590	...	1.6	0.23
1967-68 ^c		, 10mm+	1968.0	6- 7 (7.0)	3196	...	3.9	0.57
1960-61 ^d	Resistencia Province	Sample survey	Rare	RT25 1TU, 10mm+	1961.0	5- 9 (7.5)	1259	49	3.89	0.53
<u>Brazil - Bresil</u>										
1983 ^a	Rio Grande Do Sol Province	Sample survey of school enterers	0	RT23 2TU, 10mm+	1983.5	7- 8 (7.3)	11880	333	2.80	0.39
	By Provincial zone:									
	Metropolitan				1983.5	6- 7 (7.0)	3507	...	3.88	0.56
	Southern rural						4163	...	3.17	0.46
	Northern rural						3670	...	1.31	0.19
1980 ^b	Capitals of States and Territories by Region:	All school enterers	1980.5	... (7.7)				
	North						1.95
	Northeast						1.56
	Southeast						1.17
	Sao Paulo State						0.94
	South						0.78
	East Central						0.78
1970-73 ^b	Capitals of States and Territories by Region:	All school enterers	...	RT23 2TU, 10mm+	1972.0	... (7.7)				
	North						16.9	2.38
	Northeast						14.2	1.97
	Southeast						8.3	1.1
	Sao Paulo State						1.2
	South						6.7	0.90
	East Central						7.9	1.1

III. WHO South-East Asia Region - Region de l'Asie du Sud-Est de l'OMS

Year	Area Represented	Survey design	BCG (%)	Antigen and Infection Criterion	Survey Year Midpt.	Age (midpt.)	Tested N	Infect- ted N	Preva- lence (%)	Risk/ year (%)
<u>India - Inde</u>										
1979 ^e	Dodballapur in	Sample survey	14.1	RT23 1TU, 12mm+	1979.5	0- 4 (2.5)	5203	128	2.46	0.99
1974-75 ^{de}	Bangalore District	repeated in same villages	36	RT23 1TU, 10mm+	1974.8		3805	98	2.58	1.04
1977-78 ^b	3 Subdistricts in	Repeated	...	RT23 1TU, 10mm+	1977.9	0- 4 (2.5)	1492	36	2.41	0.97
1961-63 ^a	Bangalore District	sample surveys	1.5		1962.2		7981	166	2.08	0.84
1972-73 ^f	Tumkur District in	Sample survey	...	RT23 1TU, 10mm+	1973.0	0- 4 (2.5)	4277	115	2.69	1.08
1960-61 ^{cfi}	South India	repeated in same 62 villages	...		1960.9		3788	155	4.09	1.66
1978 ^g	Kashmir Valley	Sample survey	0.2	PPD-S 3TU, 12mm+	1978.7	0- 4 (2.5)	2448	56	2.29	0.92
1972 ^h	Villages in:	Selected for altitude contrast	<5	PPD-S 5TU, 12mm+	1972.8	1- 4 (3.0)				
	Kashmir						679	...	4.0	1.4
	Kulu Valley						503	...	5.2	1.8
	Lohaghat						381	...	3.1	1.0
	Pithoragarh						290	...	4.1	1.4
	Agra						530	...	5.2	1.8
	Haryana						612	...	4.4	1.5
	Rajasthan						76	...	1.3	0.4
1968-71 ^h	Northern Chingelput	Selected for BCG trial	<5	PPD-S 5TU, 12mm+	1969.9	1- 4 (3.0)	27520	...	5.1	1.7
<u>Indonesia - Indonesie</u>										
1983 ^{aefgh}	Tangerang Regency	Sample survey	18.5	RT23 2TU, 10mm+	1983.6	7-10 (8.8)	1549	286	18.46	2.29
1978	in West Java	repeated in	22.1		1978.4	7-10 (8.8)	1649	440	26.68	3.47
1972		same schools	1.9		1972.2	7-10 (9.3)	1371	497	36.25	4.73
1986 ^{bh}	Sambas in West	Selected as	0	RT23 2TU, 10mm+	1986.8	7-10 (8.8)	3839	576	15.00	1.83
1981	Kalimantan	BCG-free area			1981.8	7-10 (8.6)	2181	310	14.21	1.77
1976					1976.9	7-10 (8.6)	1655	391	23.63	3.09
1986 ^{bh}	Padang Pariaman in	Selected as	0	RT23 2TU, 10mm+	1986.9	7-10 (8.7)	1986	472	23.77	3.07
1981	West Sumatra	BCG-free area			1981.7	7-10 (8.9)	2501	363	14.51	1.75
1976					1976.5	7-10 (8.6)	1124	193	17.17	2.17
1984 ^{gh}	Malang in East Java	Selected as	0	RT23 2TU, 10mm+	1984.5	7-10 (8.3)	1406	75	5.33	0.66
1979 ^{cgh}		BCG-free area			1979.5	7-10 (8.9)	1122	86	7.66	0.89
1985 ^{cgh}	Ogan Kamering Ilir in	Selected as	0	RT23 2TU, 10mm+	1985.8	7-10 (8.9)	4840	1309	27.05	3.48
1980	South Sumatra	BCG-free area			1980.2	7-10 (8.5)	4839	1358	28.06	3.80
1975					1975.6	7-10 (8.9)	2425	701	28.91	3.76

III. WHO South-East Asia Region - Region de l'Asie du Sud-Est de l'OMS (continued)

Year	Area Represented	Survey design	BCG (%)	Antigen and Infection Criterion	Survey Year Midpt.	Age (midpt.)	Tested N	Infect- ted N	Preva- lence (%)	Risk/ year (%)
1985 ^{cgh} 1980 1975	Gowa in South Sulawesi	Selected as BCG-free area	0	RT23 2TU, 10mm+	1985.5 1980.5 1975.8	7-10 (8.9) 7-10 (8.7) 7-10 (8.2)	4001 3573 1429	1198 1076 429	29.94 30.11 30.02	3.92 4.03 4.26
1983 ^{cgh} 1978	Stabat Langkat in North Sumatra	Selected as BCG-free area	0	RT23 2TU, 10mm+	1983.2 1978.4	7-10 (8.5) 7-10 (8.9)	1894 1125	196 165	10.35 14.67	1.28 1.77
1982 ^{cgh} 1977	Hulu Sengai Tengah in South Kalimantan	Selected as BCG-free area	0	RT23 2TU, 10mm+	1982.8 1977.3	7-10 (9.0) 7-10 (8.7)	1577 1199	424 352	26.89 29.36	3.42 3.92
1984 ^{gh} 1979 1974	Pati in Central Java	Selected as BCG-free area	0	RT23 2TU, 10mm+	1984.5 1979.5 1974.5	7-10 (8.5) 7-10 (8.9) 7-10 (8.1)	2938 2197 1070	403 381 143	13.72 17.34 13.36	1.72 2.12 1.76
1983 ^g	Langsa in D.I. Aceh	Selected as BCG-free area	0	RT23 2TU, 10mm+	1983.5	7-10 (9)	24.4	3.06
1964-65 ^{dh}	Rural area in East Java	Sample survey	...	RT23 2TU, 10mm+	1965.0	5- 9 (7.5)	1633	191	11.70	1.64
<u>Thailand - Thaïlande</u>										
1977-79 ^a	Whole country	Sample survey	39.1	...	1978.5	0-14 (7.5)	15.2	2.17
1960-64 ^{bd}	Bangkok and Chiangmai Province	Sample survey	1962.5	10-14 (12.5)	29	2.7
1954 ^c	6 localities in Chiangmai and Kanchanaburi Provinces	Selected unvaccinated areas	0	... 5TU, 10mm+	1954.9	8-12 (10.5)	1578	371	23.51	2.52

IV. WHO Eastern Mediterranean Region (and Algeria) - Region de la Mediterranee Orientale de l'OMS (et Algerie)

Year	Area Represented	Survey design	BCG (%)	Antigen and Infection Criterion	Survey Year Midpt.	Age (midpt.)	Tested N	Infect ed N	Preva- lence (%)	Risk/ year (%)
<u>Algeria - Algerie</u>										
	Whole country in 6 economic zones	School survey every 5 years		RT23 2TU, 10mm+		8 (8.5)				
1985 ^d	Zone 2 Tizi Ouzou and Tlemcen	Second survey (commencing)	67.2		1985.5		2378	55	2.31	0.27
1984 ^d	Zone 5 & 6	First survey	59.1		1984.5		2125	64	3.01	0.36
1983 ^{ab}	Zone 4 incl. Batna	(nearly complete)	68.6		1983.9		1144	51	4.46	0.54
1982 ^{ab}	Zone 3 incl. Guelma		66.3		1982.9		1067	45	4.22	0.51
1981 ^{ab}	Zone 1 incl. Mascara		63.3		1981.9		1334	71	5.32	0.64
1980 ^{ab}	Zone 2 Tizi Ouzou and Tlemcen		44.6		1980.9		1844	71	3.85	0.46
1980-84	(5 Area weighted average)				1982.8		7514	302	4.02	0.48
1981 ^{ab}	Blida	Survey repeated	71.6	IP48 10TU, 6mm+	1981.5	8 (8.5)	1117	...	6.18	0.75
1976 ^{ab}	Blida	in same schools			1976.5	6-10 (8.5)	262	...	8.4	1.03
1949-52 ^c	Whole country	Mass BCG Campaign	...	IP48 5TU, 6mm+ IP48 10TU, 6mm+ Moro Patch, 3+ papules	1951.1	8 (8.5)	110547	...	31.2	4.30
	Selected areas									
	Blida								36.5	5.20
	Tizi Ouzou								25.5	3.40
	Tlemcen								24.3	3.22
	Mascara								29.7	4.06
	Batna								35.4	5.01
	Guelma								29.8	4.08
<u>Afghanistan</u>										
1982 ^{ac}	Whole country	Sample survey	...	RT23 1TU, 8mm+	1982.5	5- 9 (7.5)	881	208	23.61	3.53
1963 ^{bc}	Kabul	Selected schools	...	RT23 1TU, 8mm+	1963.5	7-12 (10.0)	19006	4322	22.74	2.55
	Jalalabad					7-12 (10.0)	3492	1083	31.01	3.64
	Mazar-e-Sharif					7-12 (10.0)	3170	1019	32.15	3.80
	Kandahar					7-12 (10.0)	4079	1586	38.88	4.80
	Pulikumry					13-15 (14.5)	1191	384	32.24	2.65
	(5 Area weighted average)					7-15 (10.2)	30938	8394	27.13	3.06
<u>Bahrain - Bahrein</u>										
1981 ^a	Whole country	All schools	0	RT23 1TU, 6mm+	1981.4	6- 7 (7.0)	6151	86	1.40	0.20
1969 ^a	Whole country	Sample survey	...	RT23 1TU, 6mm+	1969.5	6- 7 (7.0)	897	55	6.13	0.90

IV. WHO Eastern Mediterranean Region (and Algeria) - Region de la Méditerranée Orientale de l'OMS (et Algérie) (continued)

Year	Area Represented	Survey design	BCG (%)	Antigen and Infection Criterion	Survey Year Midpt.	Age (midpt.)	Tested N	Infection N	Prevalence (%)	Risk/year (%)
<u>Kuwait - Koweit</u>										
1981 ^a	Whole country	School campaign	0	RT23 ...	1981.5	4- 6 (5.2)	22710	261	1.15	0.22
1980					1980.5	4- 6 (5.1)	16149	101	0.63	0.12
1979					1979.5	4- 6 (5.3)	22674	283	1.25	0.24
1978					1978.5	4- 6 (5.2)	20843	338	1.62	0.31
1977					1977.5	4- 6 (5.3)	17444	266	1.52	0.29
1976					1976.5	4- 6 (5.1)	9018	201	2.23	0.44
1975					1975.5	4- 5 (4.5)	7665	100	1.30	0.29
1974					1974.5	4- 5 (4.2)	6722	55	0.82	0.20
1973					1973.5	4- 5 (4.3)	6363	80	1.26	0.29
1972					1972.5	4- 5 (4.7)	2258	38	1.68	0.36
1972-81	(10 Year weighted average)				1978.4	4- 6 (5.1)	131846	1723	1.31	0.26
1962-63 ^a	Whole country	Sample survey	1963.0	5- 9 (7.5)	10.74	1.50
<u>Libya Arab Jamahiriyah - Jamhiriya Araba Libyenne</u>										
1976-77 ^a	Whole country East Libya	Sample survey	63.5	RT23 2TU, Mode=16,17	1977.0	5- 9 (7.5)	1827	35	1.92	0.26
							951	...	1.9	0.26
1959 ^{ab}	East Province Cyrenaica	Sample survey	...	RT23 1TU, Mode=18,19	1959.7	5- 9 (7.5)	361	60	16.62	2.39
1954 ^{ac}	Ben Walid	Selected area	0	RT19-20-21 5TU, Mode=18,19	1954.5	5- 9 (7.5)	188	...	24.5	3.68
<u>Pakistan</u>										
1974-78 ^{abd}	Whole country except Baluchistan	Sample survey	7.8	RT23 1TU, 10mm+	1976.5	5- 9 (7.5)	2289	298	13.02	1.84
1961-62 ^{a-d}	Karachi, Rawalpindi, and Lahore	Sample survey	42.7*	RT23 1TU, 10mm+	1962.0	5- 9 (7.5)	769	178	23.15	3.45
1949-54 ^{be}	Provincial capitals Other urban areas Rural areas (3 Area median)	Mass BCG campaign	0	RT22 5TU, 5mm+	1954.1	7-14 (11.0)	47.5	5.69
									44.5	5.21
									41.0	4.68
									44.5	5.21

IV. WHO Eastern Mediterranean Region - Region de la Méditerranée Orientale de l'OMS (continued)

<u>Year</u>	<u>Area Represented</u>	<u>Survey design</u>	<u>BCG (%)</u>	<u>Antigen and Infection Criterion</u>	<u>Survey Year Midpt.</u>	<u>Age (midpt.)</u>	<u>Tested N</u>	<u>Infect- ted N</u>	<u>Preva- lence (%)</u>	<u>Risk/ year (%)</u>
<u>Syrian Arab Republic - République Arabe Syrienne</u>										
1983 ^a	Aleppo	Survey repeated in some 1978 survey schools	14.1	RT23 2TU, 10mm+	1983.8	6 (6.5)	1586	31	1.95	0.30
1978 ^{ab}	Aleppo	Sample survey of schools	8.5	RT23 2TU, 10mm+	1978.8	6 (6.5)	1845	31	1.68	0.26
1950 ^{ad}	Aleppo	BCG school campaign	0	Moro patch, 3+ papules	1950.3	6 (6.5)	15.3	2.52
1983 ^e	Homs	Sample survey of schools	17.2	RT23 2TU, 10mm+	1983.8	6 (6.5)	1182	11	0.93	0.14
1950 ^d	Homs	Mass BCG campaign	0	Moro patch, 3+ papules	1950.6	6 (6.5)	10.4	1.68
1960 ^c	Districts of Homs and Damascus, excluding City of Damascus	Sample survey	9*	RT23 1TU, 10mm+	1960.8	5- 9 (7.5)	387	21	5.43	0.74

V. WHO Western Pacific Region - Region du Pacifique Occidental de l'OMS

Year	Area Represented	Survey design	BCG (%)	Antigen and Infection Criterion	Survey Year Midpt.	Age (midpt.)	Tested N	Infect- ted N	Preva- lence (%)	Risk/ year (%)
<u>China - Chine</u>										
1979 ^{ab}	Areas without BCG, in whole country except Taiwan	266 of the 888 clusters in the national sample	0	RT23 2TU, 6mm+	1979.5	7 (7.5)	10000*	...	7.3	1.01
<u>Malaysia - Malaisie</u>										
1976-77 ^a	Whole country	Sample survey	81-86	RT23 1TU, Mode=15	1977.0	1- 9 (5.5)	1429	29	2.03	0.37
<u>Republic of Korea - Republique de Coree</u>										
1985 ^{ab}	Whole country	Sample survey	65.9	RT23 1TU, 10mm+	1985.5	0- 4 (2.8)	1420	77	5.42	1.97
1980 ^{cd}	Whole country	Sample survey	50.0	RT23 1TU, 10mm+	1980.5	3m- 4 (2.1)	1310	...	4.89	2.36
1975 ^c	Whole country	Sample survey	48.2	RT23 1TU, 10mm+	1975.5	3m- 4 (2.1)	1871	...	4.8	2.32
1970 ^c	Whole country	Sample survey	...	RT23 1TU, 10mm+	1970.5	3m- 4 (2.1)	8.5	4.14
1965 ^{cef}	Whole country, 2 strata	Sample survey	6.9	RT23 1TU, 10mm+	1965.5	0- 4 (2.5)	10.25	4.23
	Urban stratum	at 1 per 1040					994	97	9.76	4.02
	Rural stratum	at 1 per 2220					1383	144	10.41	4.30
<u>Philippines</u>										
1981-83 ^{ab}	Whole country	Sample survey	38.9	RT23 1TU, Mode=14,15	1982.5	2m- 4 (2.1)	2038	78	3.83	1.84
<u>Samoa</u>										
1975 ^a	Main island (Upolu)	Sample survey	72.6	RT23 1TU, 10mm+	1975.6	3m- 7 (2.8)	1824	25	1.37	0.49
1966-68 ^{bc}	Whole country	BCG Campaign	...	RT23 1TU, 10mm+	1967.3	0- 9 (5.0)	1710	46	2.69	0.54
1961-63 ^b	Whole country	All persons	...	RT23 1TU, 10mm+	1962.5	0- 9 (5.0)	1284	22	1.71	0.35

Symbols: * Estimated .. Not applicable ... Not available